AD	)	

Award Number: DAMD17-02-1-0150

TITLE: Synthesis of Lipoprotein Immunostimulants for Treating

Prostate Cancer

PRINCIPAL INVESTIGATOR: Jerald C. Hinshaw, Ph.D.

CONTRACTING ORGANIZATION: University of Utah

Salt Lake City, UT 84102-1870

REPORT DATE: January 2004

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

## REPORT

# DOCUMENTATION PAGE

Form Approved OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY 2. REPORT DATE 3. REPORT TYPE AND DATES COVERED (Leave blank) January 2004 Annual (1 Jan 2003 - 31 Dec 2003) . 4. TITLE AND SUBTITLE 5. FUNDING NUMBERS Synthesis of Lipoprotein Immunostimulants for Treating | DAMD17-02-1-0150 Prostate Cancer 6. AUTHOR(S) Jerald C. Hinshaw, Ph.D. 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 8. PERFORMING ORGANIZATION REPORT NUMBER University of Utah Salt Lake City, UT 84102-1870 jerald.hinshaw@hsc.utah.edu 9. SPONSORING / MONITORING 10. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) **AGENCY REPORT NUMBER** U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012 11. SUPPLEMENTARY NOTES 12a, DISTRIBUTION / AVAILABILITY STATEMENT 12b. DISTRIBUTION CODE

#### 13. ABSTRACT (Maximum 200 Words)

Approved for Public Release; Distribution Unlimited

The prospects of stimulating a patient's own immune system as a therapeutic approach to the treatment of cancer in general, and prostate cancer in particular, is intriguing. However, thus far immunotherapeutic approaches to the treatment of cancer (including prostate cancer) in the clinical setting have not been uniformly successful. We are chemically synthesizing molecular conjugates that comprise a Toll-Like Receptor (TLR) ligand covalently linked to a prostate cancer tumor-associated antigen protein or peptide. Using this tactic, we anticipate that the TLR-ligand portion of the conjugates will stimulate dendritic cells (DCs) (as well as other TLR-expressing antigen presenting cells) to secrete immune-activating cytokines. Concomitantly the tumor antigen component of the complex will be processed and presented to activated T cells. In this way, a new and potent immune system stimulation and antigen presentation mechanism aimed at the stimulation of combined CD8+ and CD4+ T-cell responses, along with B-cell activation via the innate/adaptive immune response connection, is being developed. We have prepared TLR-4 and TLR-7 ligands and are preparing mouse immunization experiments to test the efficacy of this new approach to cancer immunotherapy.

14. SUBJECT TERMS  prostate cancer immunotherapy, Toll-like receptors, chemical			15. NUMBER OF PAGES 14	
synthesis			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT	
Unclassified	Unclassified	Unclassified	Unlimited	

## **Table of Contents**

over
= 2981
able of Contents2
troduction3
ody3
ey Research Accomplishments8
eportable Outcomes8
onclusions8
eferences9
opendices10

#### A. Introduction

In this project, we are developing a new technique for stimulating a patient's own immune system as a therapeutic approach for the treatment of cancer in general, and prostate cancer, in particular. Thus far, immunotherapeutic protocols for treating cancer (including prostate cancer) in the clinical setting have not been uniformly successful<sup>1</sup>. Many of these methods have been directed toward the stimulation of a CD8+ T-cell response in the host, but have lacked a mechanism for vigorous immune system stimulation and have not combined CD4+ T-cell responses or B-cell involvement<sup>2</sup>. Recently, techniques involving stimulation of dendritic cells (DC) with tumor antigens have begun to show promise, but the process involves isolation of DCs and in vitro stimulation, followed by re-injection into the patient. Our methodology uses direct in vivo stimulation/maturation of DCs via Toll-like receptors (TLRs)<sup>3-5</sup>. We are chemically synthesizing molecular conjugates that comprise a TLR ligand covalently linked to a prostate cancer tumor-associated antigen protein or peptide. Using this tactic, we anticipate that the TLR-ligand portion of the conjugates will stimulate DCs (as well as other TLR-expressing antigen presenting cells) to secrete immuneactivating cytokines while the tumor antigen component of the complex will be processed and presented to activated T cells. We have addressed the synthesis of conjugates based on TLR-2 and TLR-7 ligands. The prostate cancer antigen portion of the complexes currently includes an MHC Class I peptide epitope from prostate-specific membrane antigen (PSMA)<sup>6</sup> and a general MHC Class II peptide epitope (PADRE)<sup>7,8</sup>. We have included the complete prostate-specific antigen (PSA) protein linked to a TLR-7 ligand. Our conjugates represent a new and potent immune system stimulation and antigen presentation mechanism with in vivo activation of DCs aimed at the stimulation of combined CD8+ and CD4+ T-cell responses along with B-cell activation through the innate/adaptive immune response connection. The conjugates are designed to elicit a vigorous immune response to prostate cancer, and may be administered by s.c. injection, or possibly, nasal or oral routes.

#### B. Body

This section describes research accomplishments associated with the tasks outlined in the original award application for Year Two of the project.

**Task 1.** Synthesize lipopeptide carriers covalently conjugated to PSA protein (months 1-14)

Following earlier efforts, considerable additional work was carried out during the second year of this program (with graduate student Jiang Sha) attempting to prepare the recombinant PSA protein, which contains a carboxyl terminal cyteine residue for chemical attachment to our TLR ligands. We have set aside this unproductive recombinant approach. Instead we set out to derivatize the wild-type protein. The thought was to convert one or more of the eleven PSA lysine amino groups to a desired mercaptan functionality. Human PSA was obtained from Research Diagnostics, Flanders, NJ. Several experiments aimed at chemically modifying the protein with thiolation reagents were carried out (Scheme 1).

Scheme 1. Attempted thiolation of PSA protein

Clean functionalization was not obtained. We suspect that the thiol-derivatized PSA may be undergoing facile intramolecular disulfide crosslinking resulting in very high molecular weight protein mixtures. Attempts to reduce these suspected disulfide linkages with dithothreitol (DTT) or triscarboxyethylphosphine were not promising, perhaps because of unwanted concomitant reduction of one or more of the five native disulfide crosslinks in the PSA protein. These reagents are also problematic, as they would be expected to react with our bromoacetamide TLR-ligand reagents, which would require the careful removal of all traces of the reagents while maintaining the thiolated protein in the reduced state. We therefore turned to directly functionalizing the PSA lysine amino groups with bromoacetamide TLR-7 ligand 1, which was prepared last year (Scheme 2).

Scheme 2. Preparation of a TLR-7- ligand- PSA conjugate

The major product(s) isolated from this reaction by HPLC was identified by MALDI mass spectroscopy as PSA appended with 2-3 TLR-7 ligands. Although the product is a mixture of functionalized proteins, we are satisfied with this material for initial immunization experiments. Using the bromoacetamide TLR ligands (**Figure 1**) that we prepared last year,

$$C_{15}H_{31} \longrightarrow NH$$

$$C_{15}H_{31} \longrightarrow OH$$

$$C_{15}H_{31} \longrightarrow OH$$

$$NH_{2}$$

$$NH_{2}$$

Pam3-cys-Bromoacetamide-TLR-2-Ligand 2

Bromoacetamide-TLR-7-Ligand 1

Figure 1. Functionalized TLR ligands

we have also synthesized several ligand-peptide conjugates for mouse immunization experiments. Scheme 3 illustrates the general synthetic route used.

Scheme 3. Preparation of peptide conjugates

Figure 2 lists the compounds that have now been prepared using Scheme 3.

Bromoacetamide-TLR-7-Ligand 1

Figure 2. TLR-2 and TLR-7 ligand peptide conjugates

The PSMA immunostimulatory conjugates are based upon the human PSMA HLA-A2 restricted peptide epitope (ALFDIESKV)<sup>6</sup> and incorporate an aminohexyl spacer between the peptide epitope and a terminal cysteine for attachment to the TLR ligands. The PADRE conjugates comprise the general CD4+ restricted peptide epitope PADRE<sup>7,8</sup> using the identical aminohexyl functionality bonded to a terminal cyteine (aK-clohexylalanine-VAAWTLKAAa-aminohexyl-C). We also synthesized the PSMA epitope covalently linked to the PADRE peptide through a central cysteine (aK-cyclohexylalanine-VAAWTLKAAa-C-ALFDIESKV) in order to evaluate the immunogenicity of conjugates, which contain both MHC Class I and MHC Class II epitopes. The TLR-2 ligand (PAM<sub>3</sub>-cys) lipopeptide conjugates proved to be quite insoluble, and thus far we have been unable to either adequately purify the materials or obtain mass spectral characterization data using MALDI, ESI or FAB ionization techniques. Similar difficulties have been noted recently for other PAM<sub>3</sub>-cys-peptide conjugates<sup>9</sup>. For this reason, we are using the TLR-7-ligand conjugates of Figure 2 in our immunization experiments.

**Task 2** Immunize HLA-DR4 and HLA-A2 transgenic mice with lipopeptide-PSA conjugates. Analyze immune responses and test responses against prostate cancer cell-lines (months 9-25)

Having prepared immunostimulating conjugates, we have begun mouse immunization experiments. A transgenic mouse colony has been established. Prostate cancer cell (LNCaP) cultures have been produced.

Founder homozygous mice carrying the HLA-A2.1 Enge transgene that express the human class MHC Class I antigen HLA-A2.1 on cells from the spleen, bone marrow and thymus were obtained from Jackson laboratories, Bar Harbor, ME (Stock Number: 003475). This surface protein recognizes the PSMA peptide epitope that is incorporated into our conjugates. A breeding colony of these mice has been established at The University of Utah. Since we have included the generalized human and mouse MHC Class II antigenic peptide epitope PADRE in our conjugate evaluation, it was not necessary to utilize transgenic mice expressing human HLA-DR4 as originally proposed. The PADRE peptide is expected to provide the desired CD4+ helper T-cell stimulation. Mice in groups of three were immunized (s.c., axilla) with the TLR-7 ligand PSMA-PADRE conjugate (Figure 2). Three levels of conjugate were examined (0.1, 1.0, and 10 μg). Mice were also immunized with the unconjugated PSMA, PADRE, and PSMA-PADRE peptides. Syngenetic C57BL/6J mice that do not express the HLA-A2 gene were used as controls. Twenty-eight days after the initial immunization, each group of mice received a boost vaccination. Eleven days post-boost, splenocytes and peripheral lymph node (PLN) lymphocytes were harvested and [<sup>3</sup>H]-thymidine cell proliferation assays were performed. Each cell population was administered [3H]-thymidine, stimulated with antigenic peptides (10, 15, 7.5, and 3.8 µg/ml) and controls, and incubated for 24 h. Radioactivity incorporation compared to controls was taken as an indication of the expansion of the cell population in response to antigenic peptide stimulation. Figure 3 indicates, as expected, that PLN lymphocytes from C57BL/6J control mice are not stimulated by the antigenic peptides. Mice immunized with 0.1 µg of the TLR-7-ligand PSMA-PADRE conjugate (Figure 4) appear to have mounted an immune response,




Figure 3. Non-transgenic control mice: No immune stimulation




Figure 4. Transgenic mice, 0.1 µg vaccinations: Induction of an immune response




Figure 5. Transgenic mice, 10 µg vaccinations: Induction of tolerance

although apparently mainly to the PADRE epitope. Based upon thymidine incorporation levels, the TKL-7 conjugate induces an approximately 2.5X increase in response over that elicited from immunization with the PSMA-PADRE peptide alone. Interestingly, vaccination with 10 µg of the conjugate (Figure 5) appears to have resulted in tolerance to the antigens. Similar, although less clear-cut, results were obtained from the splenocyte assays (data not shown). All the data are being further refined.

Preliminary chromium release experiments evaluating cytotoxic T lymphocyte killing of LNCaP prostate cancer cells (which express the PSMA protein) in culture have not yet yielded any results. In preparation for these experiments, we have experienced difficulty with the uptake (or retention) of <sup>51</sup>Cr by the LNCaP cells using standard labeling protocols. This technical problem should be solved shortly. Also, additional mouse immunization experiments with the TLR-7 conjugates in Figure 2 are underway.

Task 3. Prepare a PSMA covalent conjugate having a superior lipopeptide carrier determined from Task 2. Perform mouse immunization experiments. Test responses against prostate cancer cell-lines (months 20-36)

In view of our difficulties with the PSA protein conjugates, we have replaced the PSMA protein-conjugate work with relevant PSMA peptide epitope experiments. These conjugates have been synthesized (Figure 2) and their evaluation has been incorporated into **Task 2** and is underway.

## C. Key Research Accomplishments

This section provides a list of key accomplishments in the second year of this research.

- Functionalized ligands recognized by TLR-2 and TLR-7 have been prepared. These ligands were chemically modified with thiol-reactive haloacetamide functionality for attachment of cysteine-containing antigenic proteins and peptides.
- TLR-7 and TLR-2 ligand-antigenic peptide conjugates have been prepared as well as a TLR-7-ligand PSA protein conjugate.
- Mouse immunization experiments have begun. Initial data indicate an immunogenic response to the antigenic epitopes.

## D. Reportable Outcomes

This program supports graduate research assistant, Mr. Jiang Sha, and the results from his research will be incorporated into his dissertation.

#### E. Conclusions

Research on this program thus far has provided modified TLR-2 and TLR-7 ligands suitable

for chemical attachment of prostate cancer associated antigenic peptides and proteins. The resulting immunostimulating conjugates are being evaluated in mouse immunization experiments in order to ascertain the production of CD8+ and CD4+ T-cells as well as antibodies specific to prostate cancer antigens.

This research is significant in that it represents the first attempt to generate a host immune response to prostate cancer antigens using the potent immune system activation that arises from the innate/adaptive response connection from targeted TLR signaling in combination with antigen presentation. Evaluation of the separate conjugates will provide new insight into the effectiveness of the immune response to the same antigen stimulated by different TLRs. Such effects are not yet known. Furthermore, the possibility of an increased immune response to cancer antigens presented simultaneously with more than one TLR receptor activation path has not been explored. With our newly-synthesized conjugates, we are in a position to initiate these exciting studies.

#### F. References

- (1) Parmiani, G., Castelli, C., Dalerba, P., Mortarini, R., Rivoltini, L., Marincola, F., and Anichini, A. (2002) Cancer immunotherapy with peptide-based vaccines: What have we achieved? Where are we going? *J. Natl. Cancer Inst.* 94, 805-818.
- (2) Stevanovic, S. (2002) Identification of tumor-associated t-cell epitopes for vaccine development. *Nat. Rev. Cancer* 2, 1-7.
- (3) Kaisho, T., and Akira, A. (2002) Toll-like receptors as adjuvant receptors. *Biochim. Biophys. Acta 1589*, 1-13.
- (4) Beg, A. (2002) Endogenous ligands of toll-like receptors: Implications for regulating inflammatory and immune responses. *Trends Immunol*. 23, 509-512.
- (5) Medzhitov, R. (2001) Toll-like receptors and innate immunity. *Nat. Rev. Immunol 1*, 135-145.
- (6) Lu, J., and Celis, E. (2002) Recognition of prostate tumor cells by cytotoxic t lymphocytes specific for prostate-specific membrane antigen. *Cancer Res.* 62, 5807-5812.
- (7) Alexander, J., Sidney, J., Southwood, S., Ruppert, J., Oseroff, C., Maewal, A., Snoke, K., Serra, H., Kubo, R., Sette, A., and Grey, H. (1994) Development of high potency universal drrestricted helper epitopes by modification of high affinity dr-blocking peptides. *Immunity 1*, 751-761.
- (8) Alexander, J., Fikes, J., Hoffman, S., Franke, E., Sacci, J., Appella, E., Chisari, F., Guidotti, L., Chesnut, R., Livingston, B., and Sette, A. (1998) The optimization of helper t lymphocyte (htl) function in vaccine development. *Immunologic Res.* 18, 79-92.
- (9) Zeng, W., Ghosh, S., Lau, Y., Brown, L., and Jackson, D. (2002) Highly immunogenic and totally synthetic lipopeptides as self-adjuvanting immunocontraceptive vaccines. *J. Immunol.* 169,

## G. Appendix

Biosketches

Jerald C. Hinshaw, Principal Investigator

Jiang Sha, Graduate Research Assistant

BIOGRAPHICAL SKETCH					
NAME HINSHAW, JERALD CLYDE			POSITION TITLE Research Associate Professor		
EDUCATION/TRAINING (Begin with baccalaureate or other initial professional education, such as nursing. Include postdoctoral training.)					
INSTITUTION AND LOCATION	DEGRE	Ε	YEAR(s)	FIELD OF STUDY	
Oregon State University, Corvallis, Oregon	BS		1962 - 1966	Chemistry	
The University of Utah, Salt Lake City, Utah	PhD		1966 - 1970	Organic Chemistry	

RESEARCH AND PROFESSIONAL EXPERIENCE: DO NOT EXCEED TWO PAGES.

## **Research and Professional Experience:**

1970-1978	Advanced from Senior Research Chemist to Research Associate, Organic Research
1070 1004	Laboratory, Chemistry Division, Research Laboratories, Eastman Kodak Company
1978-1984	Scientist, Research and Development Laboratories, Thiokol Corporation
1980, 1986	Member, Utah Award Committee, Salt Lake Section, American Chemical Society
1981	Visiting Research Associate, University of Utah.
1981-1983	Chairman-Elect, Chairman, Past-Chairman, Salt Lake Section, American
	Chemical Society
1984-1990	Supervisor, Propellant Research Section, Research and Development Laboratories,
270.2770	Thiokol Corporation
1990-1999	Manager, Energetic Materials Research Department, Research and Development
	Laboratories, Thiokol Propulsion, Brigham City, Utah.
1996-1999	Member, State Advisory Council on Science and Technology (State of Utah,
2220 2222	Governor appointment)
1997,1998	Member, Utah State Governor's Medal for Excellence in Science and Technology
,	Award Committee
1997-1999	Chairman, State Advisory Council on Science and Technology (State of Utah,
200. 2000	Governor appointment)
1997-1999	Member, Utah Centers of Excellence Program Advisory Council (State of Utah,
1///-1///	Governor appointment)
2/00 7/00	
2/99-7/99	Senior Staff to the Technical Director, Science and Engineering, Thiokol Propulsion,
= 100 44 104	Brigham City, Utah
7/99-11/01	Research Assistant Professor, Department of Medicinal Chemistry, The University
	of Utah, Salt Lake City, Utah
11/01-current	Research Associate Professor, Department of Medicinal Chemistry, The University of
	Utah, Salt Lake City, Utah

## **Research Interests:**

Synthetic chemistry

Synthesis of bacterial oxidosqualene cyclase inhibitors

Cancer immunotherapy

Targeted drugs

Design and synthesis of small molecule inhibitors of protein-protein signaling

Design and synthesis of fluorescent phosphoinositide probes

Research and technology management.

#### **Honors:**

Listed in "American Men and Women of Science" Listed in "Who's Who in Technology" Named Outstanding Senior in Chemistry, 1966

National Defense Education Act Title IV Fellow, 1968-1970

Franklin Award, Thiokol Corporation recognition for outstanding technical achievement, 1995

## Publications/Patents: J. C. Hinshaw has over 50 publications and patents. A few are listed.

- P. Y. Lum, C. D. Armour, S. B. Stepaniants, G. Cavet, A. Leonardson, P. Garrett-Engele, M. K. Wolf, L. Butler, C. M. Rush, M. Bard, J. C. Hinshaw, P. Garnier, G. D. Prestwich, G. Schimmack, J. W. Phillips, C. J. Roberts, and D. D. Shoemaker, "Discovering Novel Modes of Action for Therapeutic Compounds using a Genome-wide Screen of Yeast Heterozygotes," *Cell*, 2004, 116, 121-137.
- J. C. Hinshaw, D.-Y. Suh, P. Garnier, F. S. Buckner, R. T. Eastman, S. P. T. Matsuda, B. M. Joubert, I. Coppens, K. A. Joiner, S. Merali, T. E. Nash, and G. D. Prestwich, "Oxidosqualene Cyclase Inhibitors As Antimicrobial Agents," *J. Med. Chem.*, 2003, **46**, 4240.
- T. M. McIntyre, A. V. Pontsler, A. R. Silva, A. St. Hilaire, Y. Xu, J. C. Hinshaw, G. A. Zimmerman, K. Hama, J. Aoki, H. Arai, and G. D. Prestwich, Identification of an intracellular receptor for lysophosphatidic acid (LPA): LPA is a transcellular PPARγ agonist, *Proc. Natl. Acad. Sci. USA*, 2003, **100**, 131.
- S. Davies, A. Pontsler, G. Marathe, K. Harrison, R. Murphy, J. C. Hinshaw, G. D. Prestwich, A. Hilaire, S. Prescott, G. Zimmerman, and T. McIntyre, "Oxidized Alkyl Phospholipids are Specific, High Affinity PPARg Ligands", *J. Biol. Chem.*, 2001, **276**, 16015.
- High Affinity PPARg Ligands", J. Biol. Chem., 2001, 276, 16015.
  J. C. Hinshaw and G. D. Prestwich, "The Design, Synthesis, and Evaluation of Molecules That Enable or Enhance Cellular Uptake: Peptoid Molecular Transporters", ChemTracts, Organic Chemistry, 2001, 14, 391. Commentary on the research by P. Wender, D. Mitchell, K. Pattabiraman, E. Pelkey, L. Steinman, and J. Rothbard, Proc. Natl. Acad. Sci. USA, 2000, 97, 13003.
- J. C. Hinshaw and G. D. Prestwich, "Pursuit of Optimal Carbohydrate-Based Anti-cancer Vaccines: Preparation of a Multi-antigenic Unimolecular Glycopeptide Containing the Tn, MBr1, and Lewisy Antigens", ChemTracts, Organic Chemistry, 2001, 14, 217. Commentary on the research by J. Allen, C. Harris, and S. Danishefsky, *J. Amer. Chem. Soc.*, 2001, 123, 1890.
- J. C. Hinshaw, D. W. Doll, R. J. Blau, G. K. Lund, "Metal Complexes for Use as Gas Generants," U.S. Patent 6,241,281, issued June 5, 2001.
- R.B. Wardle, R.M. Hajik, J.C. Hinshaw, T.K. Highsmith, "Process for the Large-Scale Synthesis of 4,10-Dinitro-2,6,8,12-Tetraoxa-4,10-Diazatetracyclo[5.5.0.0<sup>5,9</sup>0<sup>3,11</sup>]dodecane," U.S. Patent 6,107,483, issued August 22, 2000.
- J. C. Hinshaw, D. W. Doll, R. J. Blau, G. K. Lund, "Metal Complexes for Use as Gas Generants," U.S. Patent 6,039,820, issued March 21, 2000.
- G. D. Prestwich, F. S. Buckner, J. C. Hinshaw, "Methods Related to Steroid Metabolism of Parasites and Mycobacteria, and Treatment of Parasite and Mycobacterial Infections with an Oxidosqualene Cyclase Inhibitor", U.S. Patent Application filed June 16, 2000.
- R.B. Wardle, R.M. Hajik, J.C. Hinshaw, T.K. Highsmith, "Process for the Large-Scale Synthesis of 4,10-Dinitro-2,6,8,12-Tetraoxa-4,10-Diazatetracyclo[5.5.0.0]dodecane, "U.S. Patent 6,107,483, issued August 22, 2000.
- J. C. Hinshaw, D. W. Doll, R. J. Blau, G. K. Lund, "Metal Complexes for Use as Gas Generants," U.S. Patent 6,039,820, issued March 21, 2000.
- J. C. Hinshaw, D. W. Doll, R. J. Blau, G. K. Lund, "Metal Hydrazine Complexes Used as Gas Generants," U.S. Patent 5,970,703, issued October 26, 1999.
- J. C. Hinshaw, D. W. Doll, R. J. Blau, G. K. Lund, "Using Metal Complex Compositions As Gas Generants," U.S. Patent 5,735,118, issued April 7, 1998.
- R. R. Bennett, J. C. Hinshaw, and M. Barnes, "The Effects of Chemical Propulsion on the Environment," *Acta Astronautica*, 26, 531 (1992).
- J. C. Hinshaw and W. W. Edwards, "Synthesis of Tetranitropyrrole," J. Hetercyclic Chem., 29, 1721 (1992).

#### **BIOGRAPHICAL SKETCH**

Provide the following information for the key personnel in the order listed on Form Page 2.

NAME	POSITION TITLE				
SHA, JIANG	Graduate Research Assistant				
EDUCATION/TRAINING (Begin with baccalaureate or other initial professional education, such as nursing, and include postdoctoral training.)					
INSTITUTION AND LOCATION	DEGREE (if applicable)	YEAR(s)	FIELD OF STUDY		
Peking University, Beijing, China	B.S.	1997-2001	Biology		

RESEARCH AND PROFESSIONAL EXPERIENCE: Concluding with present position, list, in chronological order, previous employment, experience, and honors. Include present membership on any Federal Government public advisory committee. List, in chronological order, the titles, all authors, and complete references to all publications during the past three years and to representative earlier publications pertinent to this application. If the list of publications in the last three years exceeds two pages, select the most pertinent publications. **DO NOT EXCEED TWO PAGES.** 

## **Research and Professional Experience:**

2000-2001 Institution of Biophysics, Chinese Academy of Science, Bachelors' degree research

2001-2002 Molecular Biology Program, The University of Utah, Laboratory Rotation

**2002-current** Graduate Research Assistant, Department of Medicinal Chemistry,

The University of Utah, Salt Lake City